6. (a) Express the Hermitian Matrix:

$$A = \begin{bmatrix} 1 & -i & 1+i \\ i & 0 & 2-3i \\ 1-i & 2+3i & 2 \end{bmatrix}$$

as P+iQ where P is a real symmetric and Q is a real skew symmetric matrix.

(b) Using elementary row transformations, find the inverse the following matrix:

$$A = \begin{bmatrix} 2 & 3 & 4 \\ 4 & 3 & 1 \\ 1 & 2 & 4 \end{bmatrix}$$

(c) State and verify Cayley-Hamilton theorem for the following matrix:

$$\mathbf{A} = \begin{bmatrix} 2 & -1 & 1 \\ -1 & 2 & -1 \\ 1 & -1 & 2 \end{bmatrix}$$

- 7. (a) Find the mass of a plate which is formed by the co-ordinate planes and the plane $\frac{x}{a} + \frac{y}{b} + \frac{z}{c} = 1$, the density is given by $\rho = k$ xyz.
 - (b) Using Beta and Gamma functions, evaluate $\int_0^\infty \frac{dx}{1+x^4}$.
 - (c) Evaluate the integral $\int_{0}^{a} \int_{0}^{\sqrt{a^2 y^2}} (x^2 + y^2) dx dy$ by changing into polar coordinates.

(Following Paper ID and Roll No. to be filled in your Answer Book)							
PAPER ID: 9601	Roll No.			$\dot{\square}$			T

B. Tech.

(SEM. I) THEORY EXAMINATION 2011-12 MATHEMATICS—I

Time: 3 Hours

Total Marks: 100

SECTION-A

- 1. All parts of this question are compulsory: (2×10=20)
 - (a) Find the n^{th} derivative of $x^{n-1} \log x$.
 - (b) Find the Taylor's series expansion of:

$$f(xy) = x^3 + xy^2$$
 about point (2, 1).

(c) If $u = e^x \sin y$ and $v = e^x \cos y$, evaluate:

$$\frac{\partial(\mathbf{u},\mathbf{v})}{\partial(\mathbf{x},\mathbf{y})}$$

- (d) Find the minimum value of $x^2 y^2 + 6x + 12 = 0$.
- (e) Find the eigen values of the matrix $\begin{bmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{bmatrix}$.
- (f) Calculate the inverse of the natrix:

$$\begin{bmatrix} 1 & 2 \\ 5 & 7 \end{bmatrix}$$

- (g) Evaluate $\iiint_{0}^{1} xyz \, dx \, dy \, dz.$
- (h) Evaluate the area enclosed between the parabola $y x^2$ and the straight line y = x.
- (i) Find the magnitude of the gradient of the function $f = x y z^3$ at (1, 0, 2).
- (j) Write the statement of divergence theorem for a given vector field \vec{F} .

SECTION-B

- 2. Attempt any three parts of the following: (10×3=30)
 - (a) Find the eigen values and eigen vectors of the following matrix:

$$\begin{bmatrix} 2 & -1 & 1 \\ -1 & 2 & -1 \\ 1 & -2 & 2 \end{bmatrix}$$

- (b) If $y = a \cos(\log x) + b \sin(\log x)$. Find $(y_a)_a$.
- (c) The angles of a triangle are calculated from the sides a, b, c. If small changes δa , δb and δc are made in the sides, find δA , δB and δC where Δ is the area of the triangle and A, B, C are angles opposite to sides, a, b, c respectively. Also show that $\delta A + \delta B + \delta C = 0$.
- (d) Find the volume bounded by the elliptic paraboloids $z = x^2 + 9y^2$ and $z = 18 x^2 9y^2$.
- (e) If $\vec{A} = (x y)\hat{i} + (x + y)j$, evaluate $\oint_C \vec{A} \cdot d\vec{r}$ around the curve C consisting of $y = x^2$ and $y^2 = x$.

SECTION-C

Attempt any two parts from each question. All questions are compulsory. $(5\times2\times5=50)$

3. (a) If
$$y = tan^{-1} \left(\frac{a+x}{a-x} \right)$$
, prove that:

$$(a^2 + x^2) y_{n+2} + 2(n+1)x y_{n+1} + n(n+1) y_n = 0.$$

(1) If
$$u(x, y, z) = \log (\tan x + \tan y + \tan z)$$
, prove that

$$\sin 2x \frac{\partial u}{\partial x} + \sin 2y \frac{\partial u}{\partial y} + \sin 2z \frac{\partial u}{\partial z} = 2.$$

(c) Trace the curve:

$$r^2 = a^2 \cos 2\theta.$$

4. (a) If
$$y_1 = \frac{x_2 x_3}{x_1}$$
, $y_2 = \frac{x_3 x_1}{x_2}$ and $y_3 = \frac{x_1 x_2}{x_3}$, find the value

of
$$\frac{\partial (y_1, y_2, y_3)}{\partial (x_1, x_2, x_3)}$$
.

(b) Find the extreme values of:

$$f(x, y) = x^3 + y^3 - 3$$
 axy

- (c) If the base radius and height of a cone are measured as 4 cm and 8 cm with a possible error of 0.04 and 0.08 inches respectively, calculate the percentage (%) error in calculating volume of the cone.
- 5. (a) Define curl of a vector. Prove the following vector identity:

$$Div(\vec{\mathbf{u}} \times \vec{\mathbf{v}}) = Curl \vec{\mathbf{u}} \cdot \vec{\mathbf{v}} - Curl \vec{\mathbf{v}} \cdot \vec{\mathbf{u}} .$$

- (b) If $r = (x^2 + y^2 + z^2)^{1/2}$, evaluate $\nabla^2 (\log r)$.
- (c) Find the surface area of the plane x + 2y + 2z = 12 cut off by x = 0, y = 0 and $x^2 + y^2 = 16$.